

Widespread Hospital Variation in Supplementation of Breastfed Newborns

Laura P. Ward, MD, IBCLC

Multiple health organizations, including the American Academy of Pediatrics,¹ endorse the recommendation for exclusive breastfeeding for the first 6 months of life. In the United States, most women initiate breastfeeding, but a majority do not meet their intended goals of duration.² There is a complex interplay of factors that impact breastfeeding outcomes, which include sociodemographic variables, psychosocial issues, maternal health, and infant conditions. However, hospital practices also play a significant role in a mother's experience during the birth hospitalization and can be critical in setting the stage for breastfeeding success after discharge.²

In this issue of *Pediatrics*, Nguyen et al³ noted that in 2014, New York had the highest national rate of formula supplementation in breastfed newborns in the first 2 days of life. These authors sought to determine the factors that influence this finding. By using birth certificate data, they included 176 764 breastfed infants born in New York hospitals. Infants who were admitted to the NICU, transferred to other facilities, and/or exclusively formula fed were excluded. Although breastfeeding initiation was similar to the 2014 national rate (88% versus 81%), the authors found that approximately half of breastfed infants in their cohort received formula in the first 2 days. Furthermore, there was a higher likelihood of supplementation with increasing level of perinatal

care at the delivery hospitals. These findings persisted after adjusting for variables known to impact breastfeeding outcomes. There was significant interhospital variation in the rate of formula supplementation, ranging from rare (2.8%) to nearly universal (98.4%). This variance was widest in level 3 facilities (that are capable of providing complex care to high-risk women and their newborns) and narrowed with lower levels of care. These results challenge the long-held belief that hospital differences in exclusive breastfeeding rates are due to characteristics of the patient population served and not to individual hospital practices.

Evidence suggests that in-hospital supplementation negatively impacts longer term breastfeeding outcomes. Bunik et al⁴ examined multiple hospital practices and determined that exclusive breastfeeding at the time of discharge was the most important predictor of continued breastfeeding at 8 weeks. In addition, Chantry et al⁵ found that supplementation of breastfed infants prior to discharge increased the odds of not exclusively breastfeeding at 30 and 60 days. There are certainly circumstances in which supplementation is medically indicated during the hospital stay.⁶ However, the striking degree of hospital differences in early formula supplementation rates described in Nguyen's study² suggests unwarranted variation, or that which can't be explained by patient factors, and therefore represents an opportunity for improvement.

FREE

Division of Neonatology, Perinatal Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

DOI: <https://doi.org/10.1542/peds.2017-0946>

Accepted for publication Apr 20, 2017

Address correspondence to Laura P. Ward, MD, IBCLC, Division of Neonatology, Cincinnati Children's Hospital Medical Center, 3333 Burnet Ave, MLC 7009, Cincinnati, OH 45229-3039. E-mail: laura.ward@cchmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated she has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated she has no potential conflicts of interest to disclose.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2017-0142.

To cite: Ward LP. Widespread Hospital Variation in Supplementation of Breastfed Newborns. *Pediatrics*. 2017;140(1):e20170946

Quality-improvement methods that implement evidence-based approaches to hospital-based breastfeeding support can reduce variation in breastfeeding measures.⁷ Several national and statewide collaboratives supported by the Centers for Disease Control and Prevention have effected a measurable change in maternity care practices that support breastfeeding, as reflected by a national increase in maternity practices in infant nutrition and care scores.⁸

There are many questions raised by this article that require further exploration: What are the factors that drive this variation across facilities? What lessons can be learned from the centers with highest performance? Do hospitals with low rates of supplementation have fewer rotating staff? Do staff at lower-performing centers have different attitudes toward formula supplementation and find it more acceptable? What can level 3 facilities learn from centers with lower levels of care? Do units equipped to handle critically ill neonates have an altered perception of a “well” newborn and therefore a lower threshold to provide formula supplementation? A closer

examination of these differences could help inform future efforts.

A mother’s pathway to successful breastfeeding requires coordinated and multifaceted support throughout her pregnancy, delivery, and recovery well after hospital discharge. No single approach will eradicate all obstacles she may encounter along the way. Many disparities in breastfeeding are difficult to change, but hospital practices are a modifiable factor. Hospitals should work toward solutions that narrow gaps in inequities by optimizing maternity care practices and confronting obstacles that undermine breastfeeding success.

REFERENCES

1. Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2012;129(3). Available at: www.pediatrics.org/cgi/content/full/129/3/e827
2. Perrine CG, Scanlon KS, Li R, Odom E, Grummer-Strawn LM. Baby-friendly hospital practices and meeting exclusive breastfeeding intention. *Pediatrics*. 2012;130(1):54–60
3. Nguyen T, Dennison BA, Fan W, Xu C, Birkhead GS. Variation in formula

supplementation of breastfed newborn infants in New York hospitals. *Pediatrics*. 2017;140(1):e20170142

4. Bunik M, Shobe P, O’Connor ME, et al. Are 2 weeks of daily breastfeeding support insufficient to overcome the influences of formula? *Acad Pediatr*. 2010;10(1):21–28
5. Chantry CJ, Dewey KG, Peerson JM, Wagner EA, Nommsen-Rivers LA. In-hospital formula use increases early breastfeeding cessation among first-time mothers intending to exclusively breastfeed. *J Pediatr*. 2014;164(6):1339–1345.e5
6. Academy of Breastfeeding Medicine Protocol Committee. ABM clinical protocol #3: hospital guidelines for the use of supplementary feedings in the healthy term breastfed neonate, revised 2009. *Breastfeed Med*. 2009;4(3):175–182
7. Bartick M, Stuebe A, Shealy KR, Walker M, Grummer-Strawn LM. Closing the quality gap: promoting evidence-based breastfeeding care in the hospital. *Pediatrics*. 2009;124(4). Available at: www.pediatrics.org/cgi/content/full/124/4/e793
8. Grummer-Strawn LM, Shealy KR, Perrine CG, et al. Maternity care practices that support breastfeeding: CDC efforts to encourage quality improvement. *J Womens Health (Larchmt)*. 2013;22(2):107–112

Widespread Hospital Variation in Supplementation of Breastfed Newborns

Laura P. Ward

Pediatrics 2017;140;

DOI: 10.1542/peds.2017-0946 originally published online June 26, 2017;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/140/1/e20170946>

References

This article cites 8 articles, 4 of which you can access for free at:
<http://pediatrics.aappublications.org/content/140/1/e20170946.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Nutrition

http://classic.pediatrics.aappublications.org/cgi/collection/nutrition_sub

Breastfeeding

http://classic.pediatrics.aappublications.org/cgi/collection/breastfeeding_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<https://shop.aap.org/licensing-permissions/>

Reprints

Information about ordering reprints can be found online:
<http://classic.pediatrics.aappublications.org/content/reprints>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Widespread Hospital Variation in Supplementation of Breastfed Newborns

Laura P. Ward

Pediatrics 2017;140;

DOI: 10.1542/peds.2017-0946 originally published online June 26, 2017;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/140/1/e20170946>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

