

# Repeat Antenatal Betamethasone and Cardiometabolic Outcomes

Robert D. Cartwright, BSc,<sup>a</sup> Jane E. Harding, DPhil,<sup>a</sup> Caroline A. Crowther, MD,<sup>a,b</sup> Wayne S. Cutfield, MD,<sup>a,c</sup> Malcolm R. Battin, MD, MPH,<sup>a,d</sup> Stuart R. Dalziel, PhD,<sup>a,e</sup> Christopher J.D. McKinlay, PhD,<sup>a,f,g</sup> on behalf of the ACTORDS Follow-up Group

abstract

**BACKGROUND:** Repeat dose(s) of antenatal betamethasone are recommended for women at <32 weeks with ongoing risk of preterm birth. However, there is concern that use of repeat dose(s) in fetal growth restriction (FGR) may increase the risk of later cardiometabolic disease.

**METHODS:** We undertook secondary analysis of data from the Australasian Collaborative Trial of Repeat Doses of Corticosteroids Midchildhood Outcome Study to determine if FGR influences the effect of repeat betamethasone on growth and cardiometabolic function. At 6 to 8 years, children underwent anthropometry, dual energy x-ray absorptiometry, intravenous glucose tolerance testing, ambulatory blood pressure monitoring, and spirometry. FGR was defined as severe FGR at entry, cesarean delivery for FGR, or customized birth weight below the third centile.

**RESULTS:** Of 266 children assessed, FGR occurred in 43 of 127 (34%) exposed to repeat betamethasone and 44 of 139 (32%) exposed to placebo. There was an interaction between FGR and repeat betamethasone treatment for the effect on height (*z* score mean difference [95% confidence interval]; FGR: 0.59 [0.01 to 1.17]; non-FGR: -0.29 [-0.69 to 0.10]; *P* = .01). However, FGR did not influence the effect of repeat betamethasone on cardiometabolic function, which was similar in treatment groups, both in FGR and non-FGR subgroups.

**CONCLUSIONS:** Repeat antenatal betamethasone treatment had no adverse effects on cardiometabolic function, even in the presence of FGR. It may have a positive effect on height in FGR. Clinicians should use repeat doses of antenatal corticosteroids when indicated before preterm birth, regardless of FGR, in view of the associated neonatal benefits.



<sup>a</sup>Liggins Institute and <sup>d</sup>Department of Paediatrics, Child and Youth Health, University of Auckland, Auckland, New Zealand; <sup>b</sup>Department of Obstetrics and Gynaecology, School of Medicine, The University of Adelaide, Adelaide, Australia; <sup>c</sup>A Better Start, National Science Challenge, Auckland, New Zealand; <sup>e</sup>Newborn Services, National Women's Health, Auckland City Hospital, Auckland, New Zealand; <sup>f</sup>Children's Emergency Department, Starship Children's Health, Auckland, New Zealand; and <sup>g</sup>Kidz First Neonatal Care, Counties Manukau Health, Auckland, New Zealand

Mr Cartwright conducted the initial analyses and drafted the manuscript; Drs Harding, Crowther, Cutfield, and Battin conceptualized and designed the study, obtained funding, and contributed to the interpretation of data; Dr Dalziel conceptualized and designed the study and contributed to the interpretation of data; Dr McKinlay conceptualized and designed the study, obtained funding, conducted the initial analyses, and drafted the manuscript; and all authors revised the manuscript and approved the final manuscript as submitted.

This trial has been registered with the ACTRN Register (<http://anzctr.org.au>) (identifier ACTRN12606000318583).

**DOI:** <https://doi.org/10.1542/peds.2018-0522>

Accepted for publication Apr 20, 2018

**WHAT'S KNOWN ON THIS SUBJECT:** Repeat dose(s) of antenatal betamethasone are recommended for women at <32 weeks with ongoing risk of preterm birth, but there is concern that use of repeat dose(s) in fetal growth restriction (FGR) may increase the risk of later cardiometabolic disease.

**WHAT THIS STUDY ADDS:** Repeat antenatal betamethasone treatment had no adverse effects on cardiometabolic function in midchildhood, even in FGR, but may have a positive effect on height in FGR. Repeat dose(s) are recommended for preterm birth, when indicated, regardless of FGR.

**To cite:** Cartwright RD, Harding JE, Crowther CA, et al. Repeat Antenatal Betamethasone and Cardiometabolic Outcomes. *Pediatrics*. 2018;142(1):e20180522

Antenatal corticosteroid therapy is one of the most effective interventions for preterm birth, resulting in rapid maturation of fetal organ systems.<sup>1</sup> The Australasian Collaborative Trial of Repeat Doses of Corticosteroids (ACTORDS) revealed that neonatal benefit is greatest when antenatal corticosteroids are administered in repeat doses,<sup>2</sup> a finding supported by authors of systematic reviews<sup>3</sup> and animal studies.<sup>4,5</sup> Repeat doses can be used to reduce the incidence and severity of preterm lung disease and other serious neonatal morbidity, with absolute benefits similar to that of an initial course of antenatal corticosteroids.<sup>2</sup> Detailed follow-up of children in the ACTORDS at 2 and 6 to 8 years of age has revealed that repeat dose(s) of betamethasone given at intervals 7 days or more if there is ongoing risk of preterm birth within the next week are safe with no adverse effects on growth, neurodevelopment, behavior, learning, general health, lung function, bone mass, blood pressure, body composition, renal function, and glucose homeostasis.<sup>6-9</sup>

However, concern remains about whether it is safe to expose fetuses with growth restriction to repeat doses of corticosteroids, particularly because fetal growth restriction (FGR) is associated with an increased risk of cardiometabolic disease,<sup>10</sup> and increased exposure to maternal corticosteroids has been postulated as an underlying mechanism.<sup>11</sup> In addition, animal studies have revealed adverse effects of repeat antenatal corticosteroids on later cardiometabolic and lung function (effects that were most apparent in animals with FGR).<sup>12-15</sup> Further, in an observational study, antenatal corticosteroid exposure in early FGR was associated with an increased risk of short stature in childhood.<sup>16</sup>

No randomized trials have specifically evaluated the efficacy and safety of repeat antenatal

corticosteroid therapy in FGR. Therefore, we undertook a secondary analysis of the ACTORDS data to determine the influence of FGR on the effects of repeat doses of antenatal betamethasone on growth and cardiometabolic function in midchildhood.

## METHODS

### ACTORDS

The ACTORDS (international standard randomized controlled trial number 48656428) was a placebo-controlled, double-blind, randomized trial of repeat antenatal betamethasone treatment conducted at 23 collaborating hospitals across Australia and New Zealand.<sup>2</sup> Women with a single, twin, or triplet pregnancy were eligible if they were <32 weeks' gestation, had ongoing risk of preterm birth, and had received a single course of antenatal corticosteroids  $\geq 7$  days before trial entry. Exclusion criteria included chorioamnionitis requiring urgent delivery, labor at the second stage, mature fetal lung development, or further corticosteroid therapy that was considered essential.

A total of 982 women (1146 fetuses) were randomly assigned, via a central telephone service, to either the repeat betamethasone (Celestone Chronodose, comprising 7.8 mg betamethasone sodium phosphate and 6 mg betamethasone acetate; Schering-Plough, Sydney, Australia) or the masked saline placebo. Treatment was repeated each week if a woman was judged to be at continued risk of preterm birth, up to 32 weeks' gestation.

Repeat antenatal betamethasone therapy was associated with neonatal benefit, including clinically important reductions in respiratory distress syndrome (RDS), severe lung disease, and other combined serious morbidity.<sup>2</sup> At the 2 years' corrected age, neurodevelopment,

growth and general health were similar between groups.<sup>6</sup> Before the completion of the 2-year follow-up, midchildhood follow-up was planned because of concerns of the potential for long-term adverse effects of fetal corticosteroid exposure.

### Midchildhood Outcomes Study

All surviving children of mothers who had participated in and remained in the ACTORDS were invited to partake in a study of neurocognitive function, growth, and general health, including spirometry (EasyOne 2001; New Diagnostic Designs Technologies, Zurich, Switzerland), at 6 to 8 years' corrected age.<sup>7</sup> In addition, children who were assessed in New Zealand were also invited to participate in a more detailed study of cardiometabolic function, as previously described.<sup>8,9</sup> This included the measurement of body composition by whole-body dual energy radiograph absorptiometry (DXA), glucose and insulin function by a frequently sampled intravenous glucose tolerance test with Minimal Modeling (MinimalModel, version 6.02), renal function by estimated creatinine clearance, and 24-hour ambulatory blood pressure (AMBp) monitoring (Spacelabs 90217; Spacelabs Healthcare Co, Issaquah, WA). These assessments were conducted throughout New Zealand by a single team of investigators, blinded to treatment allocation. Children with severe disability or who could not assent to testing were excluded from the cardiometabolic studies. Assessment of body composition was limited to 2 centers (Auckland and Christchurch) so that DXA measurements could be performed on identical instruments (Lunar Prodigy; General Electric Healthcare, Madison, WI). Ethical approval was obtained from the New Zealand Multicentre Ethics Committee. Written informed consent was obtained from

caregivers at follow-up, and assent was sought from the children.

In this study, we hypothesized that exposure to repeat antenatal corticosteroids, compared with a single dose, would have an adverse effect on growth, cardiometabolic risk factors, bone mass, and lung function at 6 to 8 years' corrected age in children with FGR but not children with normal prenatal growth.

### Statistical Analysis

Children who underwent 1 or more tests of cardiometabolic function at 6 to 8 years' corrected age were included in this secondary analysis of data from the New Zealand ACTORDS Midchildhood Outcomes Study. The prespecified outcomes for this analysis were age- and sex-specific *z* scores for height, weight, and BMI; whole-body fat and lean mass, regional fat mass (android and gynoid), and limb lean mass; whole-body bone mineral content and area; fasting plasma glucose and insulin concentrations, and Minimal Model estimates of insulin sensitivity, acute insulin release, and disposition index; height-specific *z* scores for daytime and nighttime systolic and diastolic AMBP and percentage diurnal dip; plasma creatinine concentration and estimated glomerular filtration rate; and height- and sex-specific *z* scores for forced vital capacity (FVC), forced expiratory volume in 1 second (FEV1), FEV1/FVC ratio, and mean forced expiratory flow at 25% to 75% of FVC (FEF25-75).

FGR was defined a priori as 1 or more of the following: obstetric diagnosis of FGR at trial entry, cesarean delivery for FGR, or having a customized birth weight less than the third centile (version 6.7.8.3; Perinatal Institute, Birmingham, United Kingdom). Although this definition includes postrandomization factors, we felt that this was important because antenatal diagnosis of FGR substantially underrepresents the

true incidence of FGR in the preterm population.<sup>17,18</sup> Customized birth weight centiles, which are based on fetal growth curves, were used because these have been shown to improve detection of FGR.<sup>19,20</sup> Although infants exposed to repeat doses in the ACTORDS had slightly lower birth weight *z* scores, meta-analysis has revealed that repeat doses do not increase the risk of being small for gestational age.<sup>3</sup>


Analysis was performed by using SAS (version 9.4; SAS Institute, Inc, Cary, NC). Data are presented as number (percent), mean (SD), or median (interquartile range). Treatment groups were compared for the prespecified outcomes by using generalized linear models with adjustment for sex and gestational age at trial entry and clustering of children from multiple pregnancy by generalized estimating equations. Body composition data were additionally adjusted for current height. The influence of FGR on treatment effect was assessed by an interaction test. Treatment effects within subgroups (FGR and non-FGR) are reported as either mean difference (MD) for normally distributed data or ratio of geometric means (RGM) for positively skewed data, with 95% confidence intervals (CIs).

### RESULTS

Of the 323 surviving children eligible for follow-up in New Zealand, 308 were assessed at 6 to 8 years' corrected age, of whom 266 underwent 1 or more tests of cardiometabolic function (repeat betamethasone 127; placebo 139; Fig 1). Children who did not complete any tests of cardiometabolic function, compared with those who were assessed, had lower birth weight *z* scores, and their mothers were more likely to be multiparous but were similar for other baseline characteristics (Supplemental Table 5).

Of the 266 children included in this analysis, the rate of FGR was similar between those exposed to repeat betamethasone (34%) and placebo (32%). In the FGR subgroup, those randomly exposed to repeat betamethasone were more likely than those exposed to placebo to have received  $\geq 4$  trial treatments (repeat betamethasone 10 of 43 [27.0%]; placebo 1 of 44 [2.9%];  $P = .007$ ) and had a longer gestation (repeat betamethasone mean [SD]: 31.8 [2.7] weeks; placebo mean [SD]: 30.6 [2.7] weeks;  $P = .03$ ). Repeat betamethasone therapy reduced the severity of neonatal lung disease and the need for surfactant therapy (Table 1). In the non-FGR subgroup, there were no differences in baseline characteristics between children randomly exposed to repeat betamethasone or placebo, and, again, repeat betamethasone therapy reduced the incidence of RDS and severity of neonatal lung disease (Table 1).

The FGR subgroup, compared with non-FGR, was characterized by older maternal age (FGR mean [SD]: 32.4 [5.4] years; non-FGR mean [SD]: 30.7 [5.6] years;  $P = .03$ ), higher maternal BMI (FGR mean [SD]: 28.7 [6.9]; non-FGR mean [SD]: 25.8 [5.7];  $P = .004$ ), later gestational age at trial entry (FGR mean [SD]: 28.9 [2.0] weeks; non-FGR mean [SD]: 28.2 [2.3] weeks;  $P = .02$ ), and increased rates of multiple pregnancy (FGR: 25 of 72 [35.0%]; non-FGR: 28 of 159 [18.0%];  $P = .007$ ) and preeclampsia (FGR: 31 of 72 [43.0%]; non-FGR: 18 of 159 [11.0%];  $P \leq .001$ ) (Table 1). The FGR subgroup, compared with non-FGR, had a shorter gestation (FGR mean [SD]: 31.2 [2.7] weeks; non-FGR mean [SD]: 32.1 [3.9] weeks;  $P = .03$ ), lower birth weight *z* score (FGR mean [SD]:  $-1.1$  [0.8]; non-FGR mean [SD]: 0.1 [0.8];  $P \leq .001$ ), and an increased rate of serious neonatal morbidity (FGR: 28 of 87 [32.0%]; non-FGR: 36 of 179 [20.0%];  $P = .03$ ) (Table 1).



**FIGURE 1** Profile of participants assessed for cardiometabolic function at midchildhood. IVGTT, intravenous glucose tolerance test.

For childhood height, there was a significant interaction between repeat antenatal betamethasone therapy and FGR. In the FGR subgroup, children exposed to repeat betamethasone had larger height z scores than children exposed to placebo; no significant difference was seen between treatment groups in the non-FGR subgroup (FGR height z score MD: 0.59; 95% CI: 0.01 to 1.17; non-FGR MD: -0.29; 95% CI: -0.69 to 0.10;  $P = .01$  for interaction). In post hoc

analysis, this interaction remained significant after adjustment for ethnicity and maternal height (Table 2).

For all other cardiometabolic outcomes, treatment groups were similar, both in the FGR and non-FGR subgroups, with no evidence of an interaction effect. This included fat mass (whole body, android, gynoid), lean mass (whole-body, limb), bone mass (whole body bone mineral content and bone area) (Table 2); fasting glucose and insulin

concentrations, and Minimal Model estimates for glucose and insulin kinetics (insulin sensitivity, acute insulin release, and disposition index) (Table 3); plasma creatinine concentration, estimated glomerular filtration rate, and AMBP (daytime systolic and diastolic and nighttime systolic and diastolic z scores and diurnal systolic and diastolic percentage dip) (Table 3); and lung function (FEV1, FEV1/FVC, and FEF25-75 z scores) (Table 4).

In post hoc analysis, additional adjustment for maternal smoking during pregnancy and the number of trial treatments administered did not alter results.

## DISCUSSION

In this secondary analysis of data from the New Zealand ACTORDS Midchildhood Outcomes Study, we found that among children with FGR, exposure to repeat antenatal betamethasone was associated with increased height at midchildhood by ~0.5 SD, an effect not seen in children without FGR. Exposure to repeat antenatal betamethasone did not alter any other measures of cardiometabolic function in those with and without FGR.

FGR is associated with reduced height in childhood and adulthood, both in preterm- and term-born subjects.<sup>21</sup> This was particularly evident in the placebo group, in which children with FGR were >1 SD shorter in midchildhood than children without FGR. Contrary to our hypothesis, repeat betamethasone treatment appeared to have a positive effect on midchildhood height in children with FGR. Although we cannot exclude the possibility of a type 1 error, authors of previous studies have suggested that antenatal corticosteroid treatment may influence skeletal growth.<sup>22,23</sup> In the Auckland Steroid Trial, a single course of antenatal betamethasone compared with placebo was

**TABLE 1** Characteristics of Children, Assessed for Cardiometabolic Function at Midchildhood, and Their Mothers

	FGR Subgroup			Non-FGR Subgroup		
	Total	Repeat Betamethasone Trial Intervention	Placebo Trial Intervention	Total	Repeat Betamethasone Trial Intervention	Placebo Trial Intervention
Maternal characteristics	<i>n</i> = 72	<i>n</i> = 37	<i>n</i> = 35	<i>n</i> = 159	<i>n</i> = 75	<i>n</i> = 84
Age, y	32.4 (5.4) <sup>a</sup>	32.3 (5.9)	32.5 (4.8)	30.7 (5.6)	30 (5.5)	31.3 (5.7)
Height, cm	162.9 (6.5)	163.8 (6.9)	161.8 (5.9)	164.5 (6.8)	164.4 (6.9)	164.5 (6.8)
BMI	28.7 (6.9) <sup>a</sup>	28.5 (7.4)	28.9 (6.4)	25.8 (5.7)	26.1 (5.6)	25.5 (5.9)
Parity						
0	36.1% (26)	40.5% (15)	31.4% (11)	37.1% (59)	42.7% (32)	32.1% (27)
1–3	48.6% (35)	43.2% (16)	54.3% (19)	54.1% (86)	50.7% (38)	57.1% (48)
≥4	15.3% (11)	16.2% (6)	14.3% (5)	8.8% (14)	6.7% (5)	10.7% (9)
Multiple pregnancy	34.7% (25) <sup>a</sup>	27.0% (10)	42.9% (15)	17.6% (28)	16.0% (12)	19.0% (16)
Smoking during pregnancy	22.2% (16)	27.0% (10)	17.1% (6)	34.0% (54)	28.0% (21)	39.3% (33)
Gestational age at trial entry, wk	28.9 (2) <sup>a</sup>	28.9 (1.9)	28.9 (2.1)	28.2 (2.3)	28.2 (2.3)	28.2 (2.3)
Main reasons for risk of preterm birth <sup>b</sup>						
Preterm prelabor rupture of membranes	13.9% (10)	16.2% (6)	11.4% (4)	35.2% (56)	33.3% (25)	36.9% (31)
Preterm labor	6.9% (5) <sup>a</sup>	8.1% (3)	5.7% (2)	20.8% (33)	24.0% (18)	17.9% (15)
Severe FGR	36.1% (26) <sup>a</sup>	35.1% (13)	37.1% (13)	0.0% (0)	0.0% (0)	0.0% (0)
Preeclampsia	43.1% (31) <sup>a</sup>	40.5% (15)	45.7% (16)	11.3% (18)	12.0% (9)	10.7% (9)
Cervical incompetence	4.2% (3)	2.7% (1)	5.7% (2)	8.8% (14)	10.7% (8)	7.1% (6)
Antepartum hemorrhage	5.6% (4) <sup>a</sup>	8.1% (3)	2.9% (1)	29.6% (47)	32.0% (24)	27.4% (23)
Multiple pregnancy	4.2% (3)	2.7% (1)	5.7% (2)	5.0% (8)	8.0% (6)	2.4% (2)
Other	26.4% (19) <sup>a</sup>	24.3% (9)	28.6% (10)	12.6% (20)	6.7% (5)	17.9% (15)
No. trial treatments						
1	52.8% (38)	45.9% (17)	60.0% (21)	44.0% (70)	41.3% (31)	46.4% (39)
2 or 3	31.9% (23)	27.0% (10)	37.1% (13)	39.0% (62)	41.3% (31)	36.9% (31)
≥4	15.3% (11)	27.0% (10) <sup>c</sup>	2.9% (1)	17.0% (27)	17.3% (13)	16.7% (14)
Infant baseline characteristics	<i>n</i> = 87	<i>n</i> = 43	<i>n</i> = 44	<i>n</i> = 179	<i>n</i> = 84	<i>n</i> = 95
Sex, female	44.8% (39)	51.2% (22)	38.6% (17)	44.1% (79)	44.0% (37)	44.2% (42)
Ethnicity <sup>d</sup>						
Māori	29.9% (26)	30.2% (13)	29.5% (13)	27.9% (50)	28.6% (24)	27.4% (26)
Pacific peoples	14.9% (13)	23.3% (10)	6.8% (3)	10.6% (19)	9.5% (8)	11.6% (11)
Other non-European	6.9% (6)	9.3% (4)	4.5% (2)	5.6% (10)	7.1% (6)	4.2% (4)
European	48.3% (42)	37.2% (16)	59.1% (26)	55.9% (100)	54.8% (46)	56.8% (54)
Main neonatal outcomes						
Gestational age at birth, wk	31.2 (2.7) <sup>a</sup>	31.8 (2.6) <sup>c</sup>	30.6 (2.7)	32.1 (3.9)	31.9 (4.0)	32.2 (3.8)
Birth weight, g	1354 (454) <sup>a</sup>	1450 (446)	1260 (447)	1926 (813)	1905 (887)	1945 (745)
Birth weight z score	−1.1 (0.8) <sup>a</sup>	−1.0 (0.8)	−1.1 (0.9)	0.1 (0.8)	0.1 (0.9)	0.2 (0.7)
RDS <sup>e</sup>	43.7% (38)	34.9% (15)	52.3% (23)	40.2% (72)	29.8% (25) <sup>c</sup>	49.5% (47)
Severity of neonatal lung disease <sup>f</sup>		<sup>c</sup>			<sup>c</sup>	
Severe	12.6% (11)	4.7% (2)	20.5% (9)	12.8% (23)	7.1% (6)	17.9% (17)
Moderate	18.4% (16)	23.3% (10)	13.6% (6)	16.8% (30)	10.7% (9)	22.1% (21)
Mild	41.4% (36)	34.9% (15)	47.7% (21)	36.9% (66)	47.6% (40)	27.4% (26)
None	27.6% (24)	37.2% (16)	18.2% (8)	33.5% (60)	34.5% (29)	32.6% (31)
Mechanical ventilation	71.3% (62)	62.8% (27)	79.5% (35)	64.2% (115)	61.9% (52)	65.3% (62)
Oxygen therapy	60.9% (53)	55.8% (24)	65.9% (29)	55.9% (100)	50.0% (42)	61.1% (58)
Surfactant	27.6% (24)	16.3% (7) <sup>c</sup>	38.6% (17)	26.3% (47)	21.4% (18)	30.5% (29)
Serious neonatal morbidity <sup>g</sup>	32.2% (28) <sup>a</sup>	25.6% (11)	38.6% (17)	20.1% (36)	14.3% (12)	25.3% (24)

Data are percent (number) and mean (SD).

<sup>a</sup> *P* < .05 for comparison between subgroups (Fisher's exact test or *t* test).

<sup>b</sup> At trial entry; not mutually exclusive.

<sup>c</sup> *P* < .05 for comparison between trial intervention groups within subgroup (Fisher's exact test or *t* test).

<sup>d</sup> Ethnicity prioritized in order of Māori, Pacific, other non-European, and European.

<sup>e</sup> Defined as clinical signs of RDS and a ground glass appearance on chest radiograph.

<sup>f</sup> Defined as mild (mean airway pressure <7 cm water or fractional inspired oxygen <0.4), moderate (mean airway pressure 7 to <10 cm water or fractional inspired oxygen 0.40–0.79), or severe (mean airway pressure ≥10 cm water or fractional inspired oxygen ≥0.80).

<sup>g</sup> Defined as 1 or more of air leak syndrome, patent ductus arteriosus, need for oxygen at 36 wk postmenstrual age, severe intraventricular hemorrhage (grade 3 or 4), periventricular leukomalacia, proven necrotizing enterocolitis, or retinopathy of prematurity.

**TABLE 2** Anthropometry and Body Composition at Midchildhood of Children Exposed to Repeat Betamethasone or Placebo

Outcome	Subgroup	Repeat Betamethasone	<i>n</i>	Placebo	<i>n</i>	Treatment Effect: MD or RGM <sup>a</sup> (95% CI)	<i>P</i>
Anthropometry, mean (SD)							
Height z score	FGR	0.16 (1.15)	42	-0.45 (1.40)	44	0.59 (0.01 to 1.17)	.01
	Non-FGR	0.28 (1.26)	84	0.57 (1.23)	95	-0.29 (-0.69 to 0.10)	
Wt z score	FGR	-0.04 (1.30)	43	-0.39 (1.73)	44	0.30 (-0.35 to 0.96)	.16
	Non-FGR	0.33 (1.39)	84	0.57 (1.12)	95	-0.23 (-0.63 to 0.17)	
BMI z score	FGR	-0.18 (1.27)	42	-0.06 (1.39)	44	-0.18 (-0.77 to 0.96)	.76
	Non-FGR	0.29 (1.28)	84	0.35 (1.10)	95	-0.06 (-0.43 to 0.30)	
DXA body composition, median (interquartile range)							
Fat mass, whole body, kg	FGR	3.65 (2.60–6.22)	29	3.34 (1.72–5.14)	31	0.82 (0.64 to 1.05) <sup>a</sup>	.22
	Non-FGR	3.84 (2.44–6.33)	62	4.09 (2.77–7.16)	63	1.05 (0.88 to 1.25) <sup>a</sup>	
Fat mass, android, kg	FGR	200.1 (175.5–402.9)	29	194.1 (118.3–492.3)	31	0.84 (0.62 to 1.13) <sup>a</sup>	.33
	Non-FGR	228.0 (156.5–399.4)	62	244.8 (170.2–578.4)	63	1.04 (0.83 to 1.30) <sup>a</sup>	
Fat mass, gynoid, kg	FGR	963.2 (678.5–1419.4)	29	830.6 (509.5–1225.9)	31	0.86 (0.71 to 1.05) <sup>a</sup>	.19
	Non-FGR	949.6 (667.2–1459.0)	62	983.1 (696.6–1592.3)	63	1.06 (0.92 to 1.21) <sup>a</sup>	
Lean mass, whole body, kg	FGR	19.3 (16.4–20.8)	29	17.4 (15.6–20.2)	31	0.95 (0.91 to 1.00) <sup>a</sup>	.27
	Non-FGR	19.1 (16.5–22.4)	62	20.5 (17.8–23.6)	63	0.99 (0.96 to 1.01) <sup>a</sup>	
Lean mass, limbs, kg	FGR	7.14 (5.94–8.59)	29	6.47 (5.35–7.81)	31	0.95 (0.89 to 1.01) <sup>a</sup>	.23
	Non-FGR	7.10 (5.84–8.61)	62	7.71 (6.40–9.37)	63	1.00 (0.95 to 1.04) <sup>a</sup>	
Bone mineral content, whole body, g	FGR	876 (762–1025)	29	791 (661–1011)	31	0.94 (0.88 to 1.01) <sup>a</sup>	.41
	Non-FGR	843 (702–1018)	62	897 (792–1142)	63	0.98 (0.94 to 1.03) <sup>a</sup>	
Bone area, whole body, cm <sup>2</sup>	FGR	1058 (934–1192)	29	969 (853–1116)	31	0.97 (0.92 to 1.01) <sup>a</sup>	.44
	Non-FGR	1039 (894–1168)	62	1085 (946–1261)	63	1.00 (0.97 to 1.03) <sup>a</sup>	

<sup>a</sup> Adjusted for gestational age at trial entry and sex. Body composition comparisons were also adjusted for height. *P* value is for the test of interaction.

associated with a greater proportion of adult stature in the lower body segment,<sup>24</sup> although, in this trial and in a similar observational study of moderate-to-late preterm birth, antenatal corticosteroid exposure was not associated with altered final height.<sup>25</sup> Nevertheless, in a cohort study of adolescents with very low birth weight, adolescents exposed to antenatal corticosteroids were ~0.3 SD taller, even after adjustment for confounding factors.<sup>26</sup>

Delayed onset of puberty, leading to a prolonged period of accelerated appendicular growth, has been proposed as 1 possible explanation for these changes,<sup>8</sup> as suggested in 1 trial.<sup>27</sup> However, with our study, we aimed to assess children before pubarche.<sup>9</sup> Premature adrenarche, which is more common in children born small for gestation age, could lead to an increase in linear growth velocity, but all children were carefully examined, and none had confirmatory signs, apart from several who had also commenced puberty (Tanner stage 1).<sup>9</sup> Childhood

obesity may also accelerate linear growth,<sup>28</sup> but, in this cohort, FGR was associated with lower rather than increased BMI and fat mass.

There is ongoing debate about the long-term health effects of catch-up growth in preterm infants. Infants with growth restriction who do not exhibit catch-up growth during the neonatal period are more likely to have permanent short stature<sup>21</sup> and subnormal cognitive function in adulthood.<sup>29</sup> Conversely, human and animal studies have revealed an association between rapid catch-up growth during infancy and increased risk of cardiometabolic disease in later life.<sup>30–34</sup> Infants with FGR are predisposed to insulin resistance and central obesity, and these changes may be exacerbated by rapid catch-up growth, possibly because of altered epigenetic profiles.<sup>35,36</sup> However, in this study, FGR did not appear to be associated with increased adiposity or insulin resistance, either in children exposed to repeat betamethasone or those exposed to placebo. In addition, in the FGR subgroup,

repeat antenatal betamethasone therapy was not associated with any adverse effects on cardiometabolic function. Thus, it seems unlikely that the catch-up in height seen with repeat betamethasone in the FGR subgroup, if true, poses any risk to cardiometabolic health. We do not know the mechanism underlying this catch-up growth, although authors of an ovine study found that repeat doses of betamethasone reduced postnatal concentrations of insulin-like growth factor binding protein 3, which may potentiate the effects of insulin-like growth factor 1 on linear growth.<sup>37</sup> Investigation of the effect of repeat antenatal corticosteroids on linear growth in meta-analysis of individual patient data, including assessment of dose-response effects, may be used to help quantify this association and interaction with FGR and elucidate possible mechanisms.<sup>38</sup> A challenging issue for this study was the definition of FGR, which is conceptually straightforward but much more difficult to identify in practice. Although subgroup analysis should strictly only involve factors

**TABLE 3** Glucose, Renal, and Vascular Function at Midchildhood of Children Exposed to Repeat Betamethasone or Placebo

Outcome	Subgroup	Repeat Betamethasone	<i>n</i>	Placebo	<i>n</i>	Treatment Effect: MD or RGM <sup>a</sup> (95% CI)	<i>P</i>
Fasting plasma concentrations							
Glucose, mmol/L	FGR	4.7 (0.4)	35	4.7 (0.4)	28	0.1 (−0.1 to 0.2)	.39
	Non-FGR	4.7 (0.3)	49	4.8 (0.3)	70	−0.1 (−0.2 to 0.0)	
Insulin, mIU/L	FGR	4.8 (3.7–7.2)	35	4.5 (3.0–6.3)	28	1.19 (0.88 to 1.59) <sup>a</sup>	.68
	Non-FGR	5.2 (3.6–7.5)	49	4.9 (3.5–6.2)	70	1.05 (0.85 to 1.30) <sup>a</sup>	
Minimal model							
Insulin sensitivity, ×10 <sup>−4</sup> min <sup>−1</sup> mIU <sup>−1</sup> L	FGR	6.82 (4.49–12.91)	35	9.43 (6.94–14.26)	28	0.79 (0.57 to 1.09) <sup>a</sup>	.68
	Non-FGR	7.99 (4.91–11.02)	49	9.67 (5.58–13.81)	70	0.87 (0.70 to 1.08) <sup>a</sup>	
Acute insulin release, mIU L <sup>−1</sup> min	FGR	341 (200–752)	35	293 (174–563)	28	1.21 (0.78 to 1.86) <sup>a</sup>	.79
	Non-FGR	278 (185–504)	49	235 (171–442)	70	1.08 (0.83 to 1.41) <sup>a</sup>	
Disposition index	FGR	2723 (1902–4402)	35	3323 (1930–5226)	28	0.96 (0.64 to 1.44) <sup>a</sup>	.88
	Non-FGR	2219 (1302–3402)	49	2262 (1487–4764)	70	0.89 (0.68 to 1.16) <sup>a</sup>	
Renal function							
Plasma creatinine concentration, μmol/L	FGR	36.9 (5.3)	35	36.4 (4.1)	29	0.5 (−2.1 to 3.1)	.31
	Non-FGR	37.4 (4.7)	54	38.4 (5.8)	73	−1.0 (−2.8 to 0.9)	
Estimated glomerular filtration rate, mL per min/1.73 m <sup>2</sup>	FGR	108.0 (17.0)	35	105.3 (13.4)	29	3.1 (−5.2 to 11.5)	.70
	Non-FGR	106.1 (12.6)	54	105.3 (15.3)	73	0.8 (−4.1 to 5.7)	
AMBP							
Daytime systolic z score	FGR	−0.51 (1.05)	36	−0.53 (1.20)	41	0.04 (−0.47 to 0.54)	.45
	Non-FGR	−0.47 (0.99)	70	−0.24 (1.28)	75	−0.22 (−0.60 to 0.16)	
Daytime diastolic z score	FGR	−0.58 (1.03)	36	−0.75 (1.05)	41	0.19 (−0.30 to 0.68)	.15
	Non-FGR	−0.76 (0.89)	70	−0.52 (1.03)	75	−0.24 (−0.56 to 0.08)	
Nighttime systolic z score	FGR	0.06 (0.93)	36	0.01 (1.11)	41	0.05 (−0.44 to 0.53)	.75
	Non-FGR	−0.20 (0.92)	70	−0.15 (1.08)	75	−0.05 (−0.38 to 0.29)	
Nighttime diastolic z score	FGR	0.27 (0.92)	36	0.22 (1.04)	41	0.04 (−0.41 to 0.49)	.61
	Non-FGR	0.03 (0.95)	70	0.12 (0.94)	75	−0.09 (−0.40 to 0.22)	
Diurnal systolic % dip	FGR	10.7 (6.9)	36	12.1 (5.9)	41	−1.2 (−4.0 to 1.6)	.80
	Non-FGR	12.5 (5.2)	70	13.4 (5.0)	75	−0.9 (−2.5 to 0.7)	
Diurnal diastolic % dip	FGR	17.5 (8.4)	36	17.8 (8.3)	41	0.0 (−4.0 to 3.9)	.83
	Non-FGR	18.4 (6.5)	70	19.1 (7.1)	75	−0.7 (−2.8 to 1.5)	

Data are mean (SD) and median (interquartile range).

<sup>a</sup> Adjusted for gestational age at trial entry and sex. *P* value is for the test of interaction.**TABLE 4** Lung Function at Midchildhood of New Zealand Children Exposed to Repeat Betamethasone or Placebo

Outcome	Subgroup	Repeat Betamethasone	<i>n</i>	Placebo	<i>n</i>	Treatment Effect: MD (95% CI)	<i>P</i>
FVC z score	FGR	−0.13 (0.78)	14	−0.26 (0.85)	8	0.39 (−0.33 to 1.10)	.32
	Non-FGR	−0.13 (0.82)	30	0.05 (0.85)	36	−0.26 (−0.66 to 0.15)	
FEV1 z score	FGR	−0.75 (0.79)	14	−1.01 (1.15)	8	0.66 (−0.18 to 1.50)	.35
	Non-FGR	−1.00 (1.00)	30	−0.77 (0.91)	36	−0.20 (−0.65 to 0.25)	
FEV1/FVC z score	FGR	−1.28 (0.67)	14	−1.41 (0.87)	8	0.41 (−0.19 to 1.02)	.81
	Non-FGR	−1.52 (1.08)	30	−1.47 (0.93)	36	−0.01 (−0.47 to 0.45)	
FEF25–75 z score	FGR	−1.34 (0.92)	14	−1.37 (1.37)	8	0.62 (−0.31 to 1.56)	.74
	Non-FGR	−1.55 (1.25)	30	−1.39 (1.18)	36	−0.10 (−0.65 to 0.45)	

Data are mean (SD). Adjusted for gestational age at trial entry and sex. *P* value is for the test of interaction.

identified before randomization, we were concerned that the rate of FGR reported at trial entry (10%) underrepresented the actual degree of FGR in this high-risk cohort and could potentially obscure the influence of FGR on outcomes. Authors of studies have estimated that <25% of infants who are small for gestational

age are identified antenatally as having a potential growth problem, even in pregnancies known to be at risk, and detection rates are even lower in those not considered at risk.<sup>39,40</sup> We took a conservative approach, defining FGR as a birth weight less than the third centile, rather than the more commonly used

10th centile. Customized rather than population centiles were used because of the strong association between preterm birth and FGR.<sup>18,41</sup> Despite this narrower definition, FGR was identified in a third of the cohort, which is consistent with other reports.<sup>18</sup> As expected, the rate of serious neonatal morbidity was 60%

higher in the FGR subgroup, although the rate of severe neonatal lung disease did not differ between infants with and without FGR.

Our findings are perhaps somewhat surprising, given the number of animal studies in which authors have linked fetal corticosteroid exposure to both reduced fetal growth and cardiometabolic disease.<sup>12,42–45</sup> In addition, animal studies have suggested that maternal betamethasone administration may exacerbate fetal cardiovascular dysfunction in FGR.<sup>12,46</sup> However, human trials have consistently demonstrated that antenatal corticosteroid treatment is safe in the long-term.<sup>1,3</sup> The reason for this discrepancy has not been fully explained but may be due to the use of physiologic doses of corticosteroids, shorter durations of exposure in clinical studies, and interspecies differences in the timing of organ development.<sup>47</sup>

Authors of some studies have suggested that antenatal corticosteroid treatment may be less effective in FGR because of increased endogenous corticosteroid production and reduced clearance of corticosteroids from the fetal compartment.<sup>48,49</sup> However, in preterm lambs with FGR induced by single uterine artery ligation, maternal antenatal betamethasone continued to induce lung structural maturation and expression of surfactant proteins, although fetuses with growth restriction had increased endogenous corticosteroid concentrations.<sup>50</sup> In a large population study of preterm infants who were small for gestational age, those exposed to a single course of antenatal corticosteroids had reduced rates of death and intraventricular hemorrhage but not respiratory morbidity.<sup>51</sup> Our study has revealed that, in FGR, repeat doses of antenatal betamethasone reduced the severity of neonatal lung disease and use of surfactant, suggesting that repeat

doses may be required for pulmonary maturation in FGR in humans.

This study has some important limitations, including the limited size of the New Zealand ACTORDS cohort and the inherent risk of bias in subgroup analysis. However, we felt that it was important to undertake this exploratory analysis given the high rates of FGR among preterm infants and the ongoing concerns around efficacy and safety of repeat corticosteroid exposure in this subgroup. In addition, the ACTORDS is the only trial of repeat antenatal corticosteroids in which researchers have undertaken detailed follow-up of participants in midchildhood, at an age when cardiometabolic parameters become predictive of later risk of chronic disease.<sup>9</sup> We did not have access to paternal height, but given that maternal height was similar in randomized treatment groups, we would not expect any imbalance in midparental height.

## CONCLUSIONS

Repeat antenatal betamethasone treatment had no adverse effects on later cardiometabolic function, even in the presence of FGR. It may have a positive effect on childhood height in FGR. Clinicians should use repeat doses of antenatal corticosteroids when indicated before preterm birth, regardless of FGR, in view of the associated neonatal benefits and absence of later adverse effects.

## ACKNOWLEDGMENT

We thank the mothers and children who participated in the ACTORDS trial and this follow-up study.

## APPENDIX: COLLABORATORS

The following ACTORDS Follow-up Steering Group members are nonauthor contributors:

Ross R. Haslam (the Women's and Children's Hospital), Lex W. Doyle (Departments of Paediatrics and

Obstetrics and Gynaecology, The Royal Women's Hospital, University of Melbourne), Peter J. Anderson (Department of Paediatrics, University of Melbourne and Department of Clinical Sciences, Murdoch Children's Research Institute), Jeffrey S. Robinson (Department of Obstetrics and Gynaecology, School of Medicine, The University of Adelaide) (including authors Caroline A. Crowther and Jane E. Harding); ACTORDS clinical trial coordinator: Pat J. Ashwood (Department of Obstetrics and Gynaecology, School of Medicine, The University of Adelaide); New Zealand ACTORDS clinical trial coordinator: Coila Bevan (Liggins Institute, University of Auckland); trial statistician: Kristyn Willson (Department of Obstetrics and Gynaecology, School of Medicine, The University of Adelaide); DXA scans on children in Christchurch were conducted by Ann Mansfield (St Georges Radiology, Christchurch Radiology Group); supervision of laboratory assays was conducted by Eric Thorstensen (Liggins Institute, University of Auckland).

## ABBREVIATIONS

ACTORDS: Australasian Collaborative Trial of Repeat Doses of Corticosteroids  
AMBp: ambulatory blood pressure  
CI: confidence interval  
DXA: dual energy radiograph absorptiometry  
FEF25-75: mean forced expiratory flow at 25% to 75% of forced vital capacity  
FEV1: forced expiratory volume in 1 second  
FGR: fetal growth restriction  
FVC: forced vital capacity  
MD: mean difference  
RDS: respiratory distress syndrome  
RGM: ratio of geometric means



**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Supported through project grants from the Australian National Health and Medical Research Council (National Health and Medical Research Council: 453633), the New Zealand Health Research Council (07/204), and the Auckland Medical Research Foundation (81537).

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Roberts D, Brown J, Medley N, Dalziel SR. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev*. 2017;3:CD004454
2. Crowther CA, Haslam RR, Hiller JE, Doyle LW, Robinson JS; Australasian Collaborative Trial of Repeat Doses of Steroids (ACTORDS) Study Group. Neonatal respiratory distress syndrome after repeat exposure to antenatal corticosteroids: a randomised controlled trial. *Lancet*. 2006;367(9526):1913–1919
3. McKinlay CJD, Crowther CA, Middleton P, Harding JE. Repeat antenatal glucocorticoids for women at risk of preterm birth: a Cochrane Systematic Review. *Am J Obstet Gynecol*. 2012;206(3):187–194
4. Ikegami M, Jobe AH, Newnham J, Polk DH, Willet KE, Sly P. Repetitive prenatal glucocorticoids improve lung function and decrease growth in preterm lambs. *Am J Respir Crit Care Med*. 1997;156(1):178–184
5. Jobe AH, Newnham J, Willet K, Sly P, Ikegami M. Fetal versus maternal and gestational age effects of repetitive antenatal glucocorticoids. *Pediatrics*. 1998;102(5):1116–1125
6. Crowther CA, Doyle LW, Haslam RR, Hiller JE, Harding JE, Robinson JS; ACTORDS Study Group. Outcomes at 2 years of age after repeat doses of antenatal corticosteroids. *N Engl J Med*. 2007;357(12):1179–1189
7. Crowther CA, Anderson PJ, McKinlay CJD, et al; ACTORDS Follow-Up Group. Mid-childhood outcomes of repeat antenatal corticosteroids: a randomized controlled trial. *Pediatrics*. 2016;138(4):e20160947
8. McKinlay CJD, Cutfield WS, Battin MR, Dalziel SR, Crowther CA, Harding JE; ACTORDS Follow-Up Group. Mid-childhood bone mass after exposure to repeat doses of antenatal glucocorticoids: a randomized trial. *Pediatrics*. 2017;139(5):e20164250
9. McKinlay CJD, Cutfield WS, Battin MR, Dalziel SR, Crowther CA, Harding JE; ACTORDS Study Group. Cardiovascular risk factors in children after repeat doses of antenatal glucocorticoids: an RCT. *Pediatrics*. 2015;135(2). Available at: [www.pediatrics.org/cgi/content/full/135/2/e405](http://www.pediatrics.org/cgi/content/full/135/2/e405)
10. Demicheva E, Crispi F. Long-term follow-up of intrauterine growth restriction: cardiovascular disorders. *Fetal Diagn Ther*. 2014;36(2):143–153
11. Edwards CR, Benediktsson R, Lindsay RS, Seckl JR. Dysfunction of placental glucocorticoid barrier: link between fetal environment and adult hypertension? *Lancet*. 1993;341(8841):355–357
12. Miller SL, Supramaniam VG, Jenkin G, Walker DW, Wallace EM. Cardiovascular responses to maternal betamethasone administration in the intrauterine growth-restricted ovine fetus. *Am J Obstet Gynecol*. 2009;201(6):613.e1–613.e8
13. Johnson JWC, Mitzner W, Beck JC, et al. Long-term effects of betamethasone on fetal development. *Am J Obstet Gynecol*. 1981;141(8):1053–1064
14. Jensen EC, Gallaher BW, Breier BH, Harding JE. The effect of a chronic maternal cortisol infusion on the late-gestation fetal sheep. *J Endocrinol*. 2002;174(1):27–36
15. Zhang J, Massmann GA, Rose JC, Figueroa JP. Differential effects of clinical doses of antenatal betamethasone on nephron endowment and glomerular filtration rate in adult sheep. *Reprod Sci*. 2010;17(2):186–195
16. Schaap AH, Wolf H, Bruinse HW, Smolders-De Haas H, Van Erbruggen I, Treffers PE. Effects of antenatal corticosteroid administration on mortality and long-term morbidity in early preterm, growth-restricted infants. *Obstet Gynecol*. 2001;97(6):954–960
17. Gardosi J, Chang A, Kalyan B, Sahota D, Symonds EM. Customised antenatal growth charts. *Lancet*. 1992;339(8788):283–287
18. Cooke RWI. Conventional birth weight standards obscure fetal growth restriction in preterm infants. *Arch Dis Child Fetal Neonatal Ed*. 2007;92(3):F189–F192
19. Mongelli M, Gardosi J. Reduction of false-positive diagnosis of fetal growth restriction by application of customized fetal growth standards. *Obstet Gynecol*. 1996;88(5):844–848
20. Gardosi J. Customized fetal growth standards: rationale and clinical application. *Semin Perinatol*. 2004;28(1):33–40
21. Karlberg J, Albertsson-Wikland K. Growth in full-term small-for-gestational-age infants: from birth to final height. *Pediatr Res*. 1995;38(5):733–739
22. Swolin-Eide D, Dahlgren J, Nilsson C, Albertsson Wikland K, Holmäng A, Ohlsson C. Affected skeletal growth but normal bone mineralization in rat offspring after prenatal dexamethasone exposure. *J Endocrinol*. 2002;174(3):411–418
23. Klaus G, Jux C, Fernandez P, Rodriguez J, Himmele R, Mehls O. Suppression of

- growth plate chondrocyte proliferation by corticosteroids. *Pediatr Nephrol*. 2000;14(7):612–615
24. Dalziel SR, Walker NK, Parag V, et al. Cardiovascular risk factors after antenatal exposure to betamethasone: 30-year follow-up of a randomised controlled trial. *Lancet*. 2005;365(9474):1856–1862
  25. Norberg H, Stålnacke J, Nordenström A, Norman M. Repeat antenatal steroid exposure and later blood pressure, arterial stiffness, and metabolic profile. *J Pediatr*. 2013;163(3):711–716
  26. Doyle LW, Ford GW, Rickards AL, et al. Antenatal corticosteroids and outcome at 14 years of age in children with birth weight less than 1501 grams. *Pediatrics*. 2000;106(1). Available at: [www.pediatrics.org/cgi/content/full/106/1/e2](http://www.pediatrics.org/cgi/content/full/106/1/e2)
  27. Smolders-de Haas H, Neuvel J, Schmand B, Treffers PE, Koppe JG, Hoeks J. Physical development and medical history of children who were treated antenatally with corticosteroids to prevent respiratory distress syndrome: a 10- to 12-year follow-up. *Pediatrics*. 1990;86(1):65–70
  28. Wells JCK, Fewtrell MS, Williams JE, Haroun D, Lawson MS, Cole TJ. Body composition in normal weight, overweight and obese children: matched case-control analyses of total and regional tissue masses, and body composition trends in relation to relative weight. *Int J Obes (Lond)*. 2006;30(10):1506–1513
  29. Lundgren EM, Cnattingius S, Jonsson B, Tuvemo T. Intellectual and psychological performance in males born small for gestational age with and without catch-up growth. *Pediatr Res*. 2001;50(1):91–96
  30. Leunissen RW, Kerkhof GF, Stijnen T, Hokken-Koelega A. Timing and tempo of first-year rapid growth in relation to cardiovascular and metabolic risk profile in early adulthood. *JAMA*. 2009;301(21):2234–2242
  31. Ibáñez L, Ong K, Dunger DB, de Zegher F. Early development of adiposity and insulin resistance after catch-up weight gain in small-for-gestational-age children. *J Clin Endocrinol Metab*. 2006;91(6):2153–2158
  32. Eriksson JG, Forsén T, Tuomilehto J, Winter PD, Osmond C, Barker DJP. Catch-up growth in childhood and death from coronary heart disease: longitudinal study. *BMJ*. 1999;318(7181):427–431
  33. Ozanne SE, Nicholas Hales C. Poor fetal growth followed by rapid postnatal catch-up growth leads to premature death. *Mech Ageing Dev*. 2005;126(8):852–854
  34. Boullu-Ciocca S, Dutour A, Guillaume V, Achard V, Oliver C, Grino M. Postnatal diet-induced obesity in rats upregulates systemic and adipose tissue glucocorticoid metabolism during development and in adulthood: its relationship with the metabolic syndrome. *Diabetes*. 2005;54(1):197–203
  35. Gluckman PD, Hanson MA. Developmental and epigenetic pathways to obesity: an evolutionary-developmental perspective. *Int J Obes (Lond)*. 2008;32(suppl 7):S62–S71
  36. Khandelwal P, Jain V, Gupta AK, Kalaivani M, Paul VK. Association of early postnatal growth trajectory with body composition in term low birth weight infants. *J Dev Orig Health Dis*. 2014;5(3):189–196
  37. Gatford KL, Owens JA, Li S, et al. Repeated betamethasone treatment of pregnant sheep programs persistent reductions in circulating IGF-I and IGF-binding proteins in progeny. *Am J Physiol Endocrinol Metab*. 2008;295(1):E170–E178
  38. Crowther CA, Aghajafari F, Askie LM, et al; PRECISE Study Group. Repeat prenatal corticosteroid prior to preterm birth: a systematic review and individual participant data meta-analysis for the PRECISE study group (prenatal repeat corticosteroid international IPD study group: assessing the effects using the best level of evidence) - study protocol. *Syst Rev*. 2012;1:12
  39. Hepburn M, Rosenberg K. An audit of the detection and management of small-for-gestational age babies. *Br J Obstet Gynaecol*. 1986;93(3):212–216
  40. Backe B, Nakling J. Effectiveness of antenatal care: a population based study. *Br J Obstet Gynaecol*. 1993;100(8):727–732
  41. Anderson NH, Sadler LC, McKinlay CJD, McCowan LME. INTERGROWTH-21st vs customized birthweight standards for identification of perinatal mortality and morbidity. *Am J Obstet Gynecol*. 2016;214(4):509.e1–509.e7
  42. Berry LM, Polk DH, Ikegami M, Jobe AH, Padbury JF, Ervin MG. Preterm newborn lamb renal and cardiovascular responses after fetal or maternal antenatal betamethasone. *Am J Physiol*. 1997;272(6 pt 2):R1972–R1979
  43. Smith LM, Ervin MG, Wada N, Ikegami M, Polk DH, Jobe AH. Antenatal glucocorticoids alter postnatal preterm lamb renal and cardiovascular responses to intravascular volume expansion. *Pediatr Res*. 2000;47(5):622–627
  44. Moss TJM, Sloboda DM, Gurrin LC, Harding R, Challis JRG, Newnham JP. Programming effects in sheep of prenatal growth restriction and glucocorticoid exposure. *Am J Physiol Regul Integr Comp Physiol*. 2001;281(3):R960–R970
  45. Simchen MJ, Alkazaleh F, Adamson SL, et al. The fetal cardiovascular response to antenatal steroids in severe early-onset intrauterine growth restriction. *Am J Obstet Gynecol*. 2004;190(2):296–304
  46. Hodges RJ, Wallace EM. Mending a growth-restricted fetal heart: should we use glucocorticoids? *J Matern Fetal Neonatal Med*. 2012;25(11):2149–2153
  47. McKinlay CJ, Dalziel SR, Harding JE. Antenatal glucocorticoids: where are we after forty years? *J Dev Orig Health Dis*. 2015;6(2):127–142
  48. Torrance HL, Derks JB, Scherjon SA, Wijnberger LD, Visser GH. Is antenatal steroid treatment effective in preterm IUGR fetuses? *Acta Obstet Gynecol Scand*. 2009;88(10):1068–1073
  49. Morrison JL, Botting KJ, Soo PS, et al. Antenatal steroids and the IUGR fetus: are exposure and physiological effects on the lung and cardiovascular system the same as in normally grown fetuses? *J Pregnancy*. 2012;2012:839656
  50. Sutherland AE, Crossley KJ, Allison BJ, Jenkin G, Wallace EM, Miller SL. The effects of intrauterine growth restriction and antenatal

glucocorticoids on ovine fetal lung development. *Pediatr Res.* 2012;71(6):689–696

51. Melamed N, Pittini A, Barrett J, et al; Canadian Neonatal Network Investigators. Antenatal

corticosteroids and outcomes of small-for-gestational-age neonates. *Obstet Gynecol.* 2016;128(5):1001–1008

## Repeat Antenatal Betamethasone and Cardiometabolic Outcomes

Robert D. Cartwright, Jane E. Harding, Caroline A. Crowther, Wayne S. Cutfield,  
Malcolm R. Battin, Stuart R. Dalziel, Christopher J.D. McKinlay and on behalf of the  
ACTORDS Follow-up Group

*Pediatrics* 2018;142;

DOI: 10.1542/peds.2018-0522 originally published online June 12, 2018;

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/142/1/e20180522">http://pediatrics.aappublications.org/content/142/1/e20180522</a>
<b>References</b>	This article cites 51 articles, 11 of which you can access for free at: <a href="http://pediatrics.aappublications.org/content/142/1/e20180522#BIBL">http://pediatrics.aappublications.org/content/142/1/e20180522#BIBL</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Fetus/Newborn Infant</b> <a href="http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub">http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub</a> <b>Neonatology</b> <a href="http://www.aappublications.org/cgi/collection/neonatology_sub">http://www.aappublications.org/cgi/collection/neonatology_sub</a> <b>Cardiology</b> <a href="http://www.aappublications.org/cgi/collection/cardiology_sub">http://www.aappublications.org/cgi/collection/cardiology_sub</a> <b>Cardiovascular Disorders</b> <a href="http://www.aappublications.org/cgi/collection/cardiovascular_disorders_sub">http://www.aappublications.org/cgi/collection/cardiovascular_disorders_sub</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.aappublications.org/site/misc/Permissions.xhtml">http://www.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://www.aappublications.org/site/misc/reprints.xhtml">http://www.aappublications.org/site/misc/reprints.xhtml</a>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Repeat Antenatal Betamethasone and Cardiometabolic Outcomes**

Robert D. Cartwright, Jane E. Harding, Caroline A. Crowther, Wayne S. Cutfield,  
Malcolm R. Battin, Stuart R. Dalziel, Christopher J.D. McKinlay and on behalf of the  
ACTORDS Follow-up Group

*Pediatrics* 2018;142;

DOI: 10.1542/peds.2018-0522 originally published online June 12, 2018;

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/142/1/e20180522>

Data Supplement at:

<http://pediatrics.aappublications.org/content/suppl/2018/06/20/peds.2018-0522.DCSupplemental>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

